

Hagedorn Inflation: Open Strings on Branes Can Drive Inflation

Steven Abel^{a,1}, Katherine Freese^{b,2}, Ian I. Kogan^{c,3}

*a Dept. of Mathematical Sciences, University of Durham, Science Laboratories,
South Rd., Durham DH1 3LE, United Kingdom*

*b Michigan Center for Theoretical Physics, University of Michigan, Ann Arbor MI
48109-1120, USA*

c Theoretical Physics, 1 Keble Rd, Oxford OX1 3NP, UK

Abstract

We demonstrate an inflationary solution to the cosmological horizon problem during the Hagedorn regime in the early universe. Here the observable universe is confined to three spatial dimensions (a three-brane) embedded in higher dimensions. The only ingredients required are open strings on D-branes at temperatures close to the string scale. No potential is required. Winding modes of the strings provide a negative pressure that can drive inflation of our observable universe. Hence the mere existence of open strings on branes in the early hot phase of the universe drives Hagedorn inflation, which can be either power law or exponential. We note the amusing fact that, in the case of stationary extra dimensions, inflationary expansion takes place only for branes of three or less dimensions.

¹S.A.Abel@durham.ac.uk

²ktfreese@umich.edu

³i.kogan@physics.ox.ac.uk

1 Introduction

Inflationary cosmology [1] was proposed as a solution to the horizon, flatness, and monopole problems of the standard Hot Big Bang scenario. Inflation requires a period of accelerated expansion, $\ddot{a} > 0$, corresponding to a superluminal expansion of the scale factor. In the standard 3+1 dimensional universe, the Friedmann Robertson Walker (FRW) equations imply $\frac{\ddot{a}}{a} = -\frac{4\pi}{3m_{pl}^2}(\rho + 3p)$. Hence accelerated expansion is provided by a negative pressure. In standard inflationary models, this negative pressure is provided by a vacuum energy (a potential) with $p = -\rho$.

We have found [2] an entirely different source of negative pressure: open strings on D-branes at temperatures close to the string scale. Although Einstein's equations in higher dimensions take a different form than the FRW equation above, negative pressure can still drive inflation. Note that there is no potential of any kind in our model; instead, open strings on branes drive the inflation.

At sufficiently high temperatures and densities fundamental strings enter a curious 'long string' Hagedorn phase [3, 4, 5, 6, 7, 8]. A classical random walk picture can be used to model the behaviour of the strings in cosmological backgrounds. The particular systems we will focus on are D-branes in the weak coupling limit [9]. In particular, we consider the scenario in which our observable universe is confined to three spatial dimensions (a three-brane) embedded in higher dimensions. We will denote the rest of the universe outside of our 3-brane as the bulk. We can separate the energy momentum tensor into two components: a localized component corresponding to the D-brane tension, and a diffuse component that spreads into the bulk corresponding to open string excitations of the brane. We find an interesting type of cosmological effect of a primordial Hagedorn phase of open strings on branes:

- *Hagedorn inflation.* The transverse 'bulk' components of the energy-momentum tensor can be negative. If all of the transverse dimensions have winding modes, this negative 'pressure' causes the brane to power law inflate along its length with a scale factor that varies as $a \sim t^{4/3}$ even in the absence of a nett cosmological constant (see eq.(19)). If there are transverse dimensions that are large (in the sense that the string modes are not space-filling in these directions), then we can find exponential inflation (see eq.(20)). *No potential is required here.* Merely the existence of open strings on D-branes drives the inflation.

We begin by discussing string thermodynamics of open strings on D-branes at high temperatures near the string scale. We derive the energy-momentum tensor required for Einstein's equations from the partition function, which in turn can be derived from the

density of states in a random walk approach. Winding modes of the strings give rise to a negative ‘pressure’ in the bulk (the directions perpendicular to the three-brane on which we live). Armed with the energy-momentum tensor in Eq.(13), we examine the resultant cosmology. We can solve Einstein’s equations with various ansätze in the presence of this negative bulk component. Our primary result is that we find Hagedorn inflation of our observable universe due to the negative pressure in the bulk.

If one assumes adiabaticity, the inflationary growth period drives down the temperature of the system; eventually the temperature drop causes the universe to leave the Hagedorn regime, and consequently inflation ends automatically. In fact our solutions are only valid for small changes in the metric corresponding to small changes in the volumes, i.e., we can demonstrate instability to inflationary expansion but cannot follow the solutions further. If one uses the solutions beyond the region of their validity, the period of superluminal growth ends too quickly to solve cosmological problems. Hence we do also discuss how, in non-adiabatic systems, inflation can be sustained.

An amusing result is that power law inflation in the universal high energy system only takes place if the number of large dimensions on the brane is $p \leq 3$. In addition, branes tend to ‘melt’ unless $p \leq 4$. Hence one can speculate on the role of these effects in the fact that our observable universe has three large dimensions.

2 The Hagedorn phase and random walks

We here derive the thermodynamic properties of D-branes using a random walk approach. The results are independent of the details of the compactification and the degree of supersymmetry, and agree with previous work [7] assuming toroidal compactification in microcanonical and canonical ensembles. The reader interested only in the result for the energy-momentum tensor and the resultant cosmology should proceed directly to Eq.(13).

2.1 String thermodynamics and the Hagedorn phase

The Hagedorn phase arises in theories containing fundamental strings because they have a large number of internal degrees of freedom. Because of the existence of many oscillator modes, the density of states grows exponentially with single string energy ε , $\omega(\varepsilon) \sim \varepsilon^{-b} e^{\beta_H \varepsilon}$, where the inverse Hagedorn temperature β_H (where $\beta = 1/T$) and the exponent b depend on the particular theory in question (for example heterotic or type II) [3]. For type I, IIA, IIB strings the numerical value of the inverse Hagedorn temperature is $\beta_H = 2\sqrt{2}\pi$ in string units. Thermodynamic quantities, such as the entropy, diverge at the Hagedorn temperature. The partition function Z is obtained as the integral of the density of states times the usual Boltzmann factor $e^{-\beta\varepsilon}$, $Z \propto \int d\varepsilon \varepsilon^{-b} e^{(\beta_H - \beta)\varepsilon}$ which

diverges at $\beta = \beta_H$ for $b \leq 2$.

In the Hagedorn regime, fundamental strings can be described as ‘long strings’ in a random walk analogy. The size of the random walk (the distance covered by the string) is given by the length scale $\sqrt{\varepsilon}$. The physical setup is shown in Figure 1.




Figure 1: We consider a volume, portrayed by the box, containing a single brane embedded in the bulk. The quantity $R_{\parallel}$ indicates the size of the dimensions parallel to the brane (for cosmology, $R_{\parallel} \rightarrow \infty$) and $R_{\perp}$ indicates the size of the dimensions perpendicular to the brane. The random walk of strings attached to the brane traverses a distance $\sqrt{\varepsilon}$. Figs. 1a represents the case $\sqrt{\varepsilon} \ll R_{\perp}$. Fig. 1b represents the case $\sqrt{\varepsilon} \gg R_{\perp}$. This latter case is our standard high energy case in which all modes are space-filling (effectively, “winding modes”).

We consider a volume, portrayed by the box, containing a single brane. The quantity $R_{\parallel}$ indicates the size of the dimensions parallel to the brane (for cosmology, $R_{\parallel} \rightarrow \infty$) and $R_{\perp}$ indicates the size of the dimensions perpendicular to the brane. The random walk of strings attached to the brane traverses a distance $\sqrt{\varepsilon}$. Figs. 1a and 1b show the cases $\sqrt{\varepsilon} \ll R_{\perp}$ and $\sqrt{\varepsilon} \gg R_{\perp}$ respectively. In this paper we are particularly interested in the high energy case of fig. 1b, in which the strings fill all the space; for the case of toroidal compactification this corresponds to winding modes. We will use the nomenclature ‘winding modes’ to encompass these space-filling modes regardless of the type of compactification. We define d_o to be the number of dimensions transverse to the brane in which there are *no windings*. As our standard ‘high-energy’ regime, we will take the case of $d_o = 0$, so that all dimensions have windings; this is the system which is always reached provided that the energy density is high enough.

Previously, [7] obtained the distribution function $\omega(\varepsilon)$ for open strings attached to a brane:

$$\omega(\varepsilon)_{\text{open}} \sim \frac{V_{\parallel}}{\varepsilon^{d_{\perp}/2}} \exp(\beta_H \varepsilon) \quad R_{\perp} \gg \sqrt{\varepsilon}. \quad (1)$$

and

$$\omega(\varepsilon)_{\text{open}} \sim \frac{V_{\parallel}}{V_{\perp}} \exp(\beta_H \varepsilon) \quad R_{\perp} \ll \sqrt{\varepsilon}. \quad (2)$$

Here, $d_{\perp}$ is the number of dimensions perpendicular to the brane, and $V_{\parallel}$ and $V_{\perp}$ are the volumes parallel and perpendicular to the brane respectively.

One can now adapt the random walk to a cosmological background. In the case of a non-trivial metric the most natural interpretation of the parameter ε is that it is the proper length of a string in the bulk and certainly we can always go to the local inertial frame in which a small portion of the string has the usual Euclidean energy $\equiv$ length equivalence.

We make the usual quasi-equilibrium approximation that equilibrium is established much more quickly than any change in the metric so that the metric may be taken to be approximately constant when evaluating properties such as density. We assume that the metric is expressed in terms of parallel dimensions x and transverse ones y

$$ds^2 = -n^2 dt^2 + g_{\parallel ij} dx^i dx^j + g_{\perp nm} dy^n dy^m, \quad (3)$$

with the brane lying at $y^n = 0$. We define an averaging over the extra dimension with an overbar,

$$\overline{O} = \frac{\int dy \sqrt{g_{\perp}} O(y)}{\int dy' \sqrt{g_{\perp}}}. \quad (4)$$

The density of states for the limiting systems is found to be the same expression as in the flat space case of eq. (2) but with all volumes averaged over transverse dimensions as in eq.(4).

2.2 Partition Function

Now that we have obtained a density of states of open strings attached to branes near the Hagedorn temperature, we can find the partition function at temperature $1/\beta$,

$$\log Z(\beta, \overline{V}_{\parallel}, \overline{V}_{\perp}) \sim \int d\varepsilon \omega(\varepsilon) e^{-\beta\varepsilon}. \quad (5)$$

In the standard $d_o = 0$ case,

$$\log Z(\beta, \overline{V}_{\parallel}, \overline{V}_{\perp}) = 2 \frac{\overline{V}_{\parallel}^2 \beta_H^2}{\overline{V}_{\perp}(\beta^2 - \beta_H^2)} + \text{nonsingular cutoff terms} \quad (6)$$

which are the successive terms in a saddle point approximation. In [2], we present the density of states and partition function for arbitrary values of d_o as well.

3 Stress-energy tensor $T_{\mu\nu}$ in a bulk Hagedorn phase

We now use these thermodynamic results to find the bulk energy momentum tensor during the Hagedorn regime. We may find the energy momentum tensor from

$$\langle T_{\nu}^{\mu} \rangle = 2 \frac{g^{\mu\rho}}{\sqrt{g}} \frac{\delta \log Z(\beta, \overline{V}_{\parallel}, \overline{V}_{\perp})}{\delta g^{\rho\nu}}. \quad (7)$$

For now, we take a single extra dimensions, so that $\mu, \nu = 0, 1, 2, 3, 5$ and 5 is the extra dimension. We will treat the functional derivative with respect to $g_{\mu\nu}$ in the following way. We assume that small changes in the metric correspond to making small changes in the volumes in $\log Z$, *e.g.* for a single extra dimension

$$\frac{\delta Z}{\delta g_{55}} = \int dx' \frac{\delta Z}{\delta \overline{V}_{\perp}(x')} \frac{\delta \overline{V}_{\perp}(x')}{\delta g_{55}}. \quad (8)$$

Then, in the case of only one extra dimension, we can write

$$\overline{V}_{\perp} = \int dy \sqrt{g_{55}} = \frac{\int d^5 x \sqrt{g_{55}}}{\int d^4 x} = \frac{1}{V_{\parallel} \beta} \int d^5 x \sqrt{g_{55}}, \quad (9)$$

and

$$\frac{\delta \overline{V}_{\perp}}{\delta g_{55}} = \frac{1}{2\sqrt{g_{55}} V_{\parallel} \beta}. \quad (10)$$

Then from eqs.(8–10) we can determine the functional derivative in eq.(7). Our ansatz automatically means that $T_{05} = 0$ and hence $G_{05} = 0$; in other words we are not considering energy exchange between the brane and the bulk. (In general there might be energy flux between the two.)

We now summarize the results for the energy-momentum tensor. We first drop the overline notation of the previous section and simply redefine $V_{\perp}$ and $V_{\parallel}$ to be the transverse and parallel volumes covariantly averaged over the region of the transverse dimensions covered by the strings. We define an energy density ρ of strings

$$\rho = \frac{E}{V_{\parallel}} \quad (11)$$

where E is the total energy of the string system. We define

$$\gamma \equiv d_o/2 - 1 \quad (12)$$

where d_o is the number of dimensions with no windings.

We find that the ‘bulk’ components of the energy momentum tensor are given by

$$\begin{aligned} \hat{T}_0^0 &= -\hat{\rho} \\ \hat{T}_i^i &= \hat{p}_{\gamma} \\ \hat{T}_m^m \equiv p_{bulk} &\approx \begin{cases} -\hat{p}_{\gamma} & \text{transverse with windings} \\ 0 & \text{transverse without windings,} \end{cases} \end{aligned} \quad (13)$$

where $\hat{\rho} = \rho/V_\perp$ and

$$\hat{p}_\gamma = \begin{cases} \frac{1}{V_\perp^{3/2}} \rho^{\frac{\gamma}{\gamma-1}} & \text{if } \gamma = -1, -\frac{1}{2}, \frac{1}{2} \\ \frac{1}{V_\perp^{3/2}} \log \rho & \text{if } \gamma = 0 \\ \frac{1}{V_\perp^{3/2}} e^{-\rho} & \text{if } \gamma = 1, \end{cases} \quad (14)$$

wherever there are strings present, and zero otherwise. In particular, for our standard high energy case of $\gamma = -1$, the negative bulk pressure is:

$$p_{bulk} \sim -\rho^{1/2}/V_\perp^{3/2}. \quad (15)$$

As expected $\hat{T}_0^0$ resembles the local energy density of strings. The $\hat{T}_i^i$ represents a relatively small pressure coming from Kaluza-Klein modes in the Neumann directions and $\hat{T}_m^m$ is a *negative* pressure coming from winding modes in the Dirichlet directions. If we T-dualize the Dirichlet directions these ‘winding modes’ also become Kaluza-Klein modes in Neumann-directions and T_m^m becomes positive. Thus negative T_m^m reflects the fact that we have T-dualized a dimension much smaller than the string scale thereby reversing the pressure. For this reason negative T_μ^μ is expected to be a general feature of space-filling excitations in transverse dimensions. The most important result of this section is the negative bulk pressure found in Eqs.(13) and (15).

4 Cosmological Equations in $D = 5$

We now consider a D-brane configuration that has 3 large parallel dimensions (*i.e.* the ‘observable universe’) and only one transverse dimension y that supports winding modes. For now, we also assume adiabaticity.

We use the metric,

$$ds^2 = -n^2 dt^2 + a^2 d\mathbf{x}^2 + b^2 dy^2, \quad (16)$$

which foliates the space into flat, homogeneous, and isotropic spatial 3-planes. Here $\mathbf{x} = x_1, x_2, x_3$ are the coordinates on the spatial 3-planes while y is the coordinate of the extra dimension. For simplicity, we make a further restriction by imposing Z_2 symmetry under $y \rightarrow -y$. Without any loss of generality we choose the 3-brane of the ‘observable universe’ to be fixed at $y = 0$. We can associate a scale factor with the parallel dimensions $a(t, y)$ and one for the ‘extra’ transverse dimensions $b(t, y)$. We define $a_0(t(\tau)) = a(t, 0)$ as the scale factor describing the expansion of the 3-brane where $t(\tau) \equiv \int d\tau n(\tau, y = 0)$ is the proper time of a comoving observer.

Several authors [10] - [12] have presented the bulk Einstein equations. For illustrative purposes we restate the results for the 55 (yy) equation:

$$\hat{G}_{55} = 3 \left\{ \frac{a'}{a} \left(\frac{a'}{a} + \frac{n'}{n} \right) - \frac{b^2}{n^2} \left[\frac{\dot{a}}{a} \left(\frac{\dot{a}}{a} - \frac{\dot{n}}{n} \right) + \frac{\ddot{a}}{a} \right] \right\} = \hat{\kappa}^2 \hat{T}_{55}, \quad (17)$$

where $\hat{\kappa}^2 = 8\pi\hat{G} = 8\pi/M_5^3$, M_5 is the five-dimensional Planck mass, and the dots and primes denote differentiation with respect to t and y , respectively. As stated earlier, our ansatz implies that $T_{05} = 0$ in the bulk. Shortly we will use the $T_{\mu\nu}$ appropriate to the Hagedorn regime on the right hand side of Einstein's equations.

From Eqn.(17), one can see right away the importance of negative components of the bulk energy momentum tensor. In the case where time derivatives of a dominate on the left hand side of the equation, one has $\frac{b^2}{n^2} \left[\left(\frac{\dot{a}}{a} \right)^2 + \frac{\ddot{a}}{a} \right] = -\hat{\kappa}^2 \hat{T}_{55}$. One can see that accelerated expansion $\ddot{a} > 0$, which is required for inflation, takes place with a negative $\hat{T}_{55}$ such as that found in Eqns.(13) and (15).

In addition to the bulk Einstein equations, we have boundary conditions (known as the Israel jump conditions) due to the fact that our observable brane is embedded in the bulk. We will assume that the energy momentum tensor on the boundary can be written in a perfect fluid form, $t_0^0 = \rho_{br}$ and $t_1^1 = -p_{br}$, where ρ_{br} and p_{br} are the energy density and pressure, respectively, measured by a comoving observer. For the metric in eq.(16) and the brane at a Z_2 symmetry fixed plane, the Israel conditions become

$$\begin{aligned} 3[a'/a]_0 &= -\hat{\kappa}^2 b_0 \rho_{br} \\ 3[n'/n]_0 &= \hat{\kappa}^2 b_0 (2\rho_{br} + 3p_{br}). \end{aligned} \quad (18)$$

Here, subscript 0 refers to quantities evaluated at the location of the brane at $y = 0$, and $[X] \equiv Xy \rightarrow 0^+ - X(y \rightarrow 0^-)$ indicates the "jump" in the quantity X as one crosses the boundary (brane) at $y = 0$ (the value of X as one approaches the brane from one side minus the value of X as one approaches it from the other side).

5 Results: Behaviour of scale factors $a(t)$ and $b(t)$: Inflation without Inflavons

Our results are obtained by solving Einstein's equations together with the constraints provided by the Israel conditions. We use the energy momentum tensor derived above in Eq.(13) appropriate to a primordial Hagedorn epoch. First we will consider the case where the brane is energy/pressureless and that $\dot{b} = 0$ (where b is the scale factor of the extra dimension). We then catalogue more general types of behaviour that can occur for $\dot{b} \neq 0$.

5.1 Case I: $\Lambda_{br} = \Lambda_{bulk} = \dot{b} = 0$

Let us first impose $\dot{b} = 0$ simply as an external condition; *i.e.* the extra dimension is fixed in time. We also set the cosmological constants both on the brane and in the bulk to zero, $\Lambda_{br} = \Lambda_{bulk} = 0$. All other contributions to the energy momentum localized on the brane (for example massless Yang-Mills degrees of freedom) have an entropy that is subdominant to the limiting bulk degrees of freedom as long as the string energy density $\rho > \rho_c$ (where ρ_c is a critical Hagedorn density of order 1 in string units). For the moment we will therefore neglect the brane energy-momentum and set $a'_0(t) = n'_0(t) = 0$. In this discussion the brane at $y = 0$ is playing no role in determining the evolution of the cosmology; the scale factor a_0 changes purely as a result of the bulk equations. It is simple to solve the 55 equation for the scale factor $a_0(t)$ on our brane (at $y = 0$).

Standard case with $d_o = 0$: In the high energy case with windings in all transverse dimensions we see from Eqs.(13) and (15) that $T_5^5 \sim \sqrt{\rho} \sim a^{-3/2}$ (where, again, ρ is the local energy density measured with respect to the volume $V_{||}$) and hence,

$$a_0(t) \sim t^{4/3}, \quad (19)$$

i.e. power law inflation. For the more general case of ‘p’ large parallel dimensions in any number of extra perpendicular dimensions, we find $T_5^5 \sim a^{-p/2}$ and $a \sim t^{4/p}$. Here adiabaticity has been assumed in taking $E \sim S \sim \text{const.}$ We note the amusing result that inflation requires $p \leq 3$ (for our brane $p = 3$) and speculate on its role in our three large extra dimensions.

Case with $d_o = 2$: In the case of 2 dimensions with no windings, we find a period of *exponential inflation*:

$$\frac{a_0(t)}{a_0(0)} = \exp \left(-\frac{t^2 \hat{\kappa}^2}{8V_{\perp}^{\frac{3}{2}}} + t \sqrt{\frac{\hat{\kappa}^2}{6V_{\perp}^{\frac{3}{2}}} \log(\rho(0)e^{3/4})} \right) \quad (20)$$

where $\rho(0)$ is the initial density at $t = 0$. Initially the second term in the exponent dominates and there is exponential inflation. This solution has an automatic end to inflation when $\rho(t) \sim 1$ (in string units), just as the system is dropping out of the Hagedorn phase.

Other systems: We summarize the energy momentum tensors and cosmological behavior of $a_0(t)$ for systems with d_o perpendicular dimensions without windings in Table 1, where $p = 3$ for our universe as a 3-brane. Superluminal expansion is found for systems with $d_o \leq 2$, *i.e.*, no more than two perpendicular dimensions without windings. In all cases, we have also found solutions to the T_{00} and T_{ii} equations.

In adiabatic systems the Hagedorn regime and hence the inflationary behaviour we have found eventually come to an end. For a system to be in the Hagedorn regime

requires an entropy density higher than the critical Hagedorn density (of order 1 in string units). Below this density the energy momentum tensor and hence the cosmology is governed by the massless relativistic Yang-Mills gas (the gas is present on the brane even in the Hagedorn regime but is subdominant). Thus there is no problem exiting from the inflationary behaviour. In fact the main issue is how long inflation can last. We return to this question later when we discuss how inflation can be sustained.

regime	$\rho(\beta) = E/V_{\parallel}$	$-p_{bulk}$	$a_0(t)/a_0(0)$
$d_o = 0$	$\frac{1}{V_{\perp}}(\beta - \beta_H)^{-2}$	$\rho^{\frac{1}{2}}$	$t^{\frac{4}{p}}$
$d_o = 1$	$(\beta - \beta_H)^{-\frac{3}{2}}$	$\rho^{\frac{1}{3}}$	$t^{\frac{6}{p}}$
$d_o = 2$	$(\beta - \beta_H)^{-1}$	$\log \rho$	$\exp(-Ct^2 + Dt)$
$d_o = 3$	$(\beta - \beta_H)^{-\frac{1}{2}}$	ρ^{-1}	$t^{-\frac{2}{p}}$
$d_o = 4$	$-\log(\beta - \beta_H)$	$e^{-\rho}$	const

Table 1: Cosmological regimes for open strings in the Hagedorn phase with p large parallel dimensions, where $p = 3$ for our observable universe as a 3-brane. Here d_o is the number of dimensions transverse to the brane in which there are no windings; in our standard “high-energy” regime we take $d_o = 0$. Here ρ is the energy density of strings, *i.e.*, the T_{00} component of the bulk energy-momentum tensor. In addition, $T_{55} = p_{bulk}$ is the negative bulk pressure that drives inflation of our scale factor $a_0(t)$. Note inflationary expansion for $d_o \leq 2$, *i.e.*, at most 2 directions without windings. The constants C and D are given for $p = 3$ in eq.(20).

5.2 Case II: $\frac{\hat{\kappa}^2}{12}\Lambda_{br}^2 - \Lambda_{bulk} = 0$ and $\dot{b} \neq 0$

We also studied the case where the extra dimension is not stabilized but there is still no nett cosmological constant. By ‘nett’ we mean that the contribution from a' and n' in G_5^5 cancels the bulk contribution on the RHS of the T_5^5 equation. The condition for this is

$$\Lambda_{nett} = -\frac{\hat{\kappa}^2}{12}\Lambda_{br}^2 + \Lambda_{bulk} = 0. \quad (21)$$

Using the 55 equation, we catalogued some possible types of behavior for a with different ansätze for b . Since $\alpha \approx 1$, for $\gamma = -1, -\frac{1}{2}, \frac{1}{2}$ ($\gamma = d_o/2 - 1$), the pressure is

$$-p_{bulk} \propto a^{\frac{-3\gamma}{\gamma-1}} b^{\frac{-3}{2}}. \quad (22)$$

Taking $\Lambda_{nett} = 0$, we find a family of power law solutions for $\gamma = -1, -\frac{1}{2}, \frac{1}{2}$ of the form

$$\begin{aligned} a_0(t) &\approx At^q \\ b_0(t) &\approx Bt^r \end{aligned} \quad (23)$$

where subscript-0 indicates values at $y = 0$,

$$q = \frac{\gamma - 1}{2\gamma} \left(\frac{4}{3} - r \right), \quad (24)$$

and r is arbitrary. For our standard $\gamma = -1$, we find superluminal solutions with $q > 1$ for $r < 1/3$. Hence one can have an inflating brane with a shrinking bulk ($r < 0$) or with a growing bulk ($0 < r < 1/3$). We also find a family of hyperbolic solutions with

$$\frac{a_o(t)}{a_o(0)} = \left(\frac{\sinh 2C(t + t_1)}{\sinh 2Ct_1} \right)^{1/2}. \quad (25)$$

These solutions can give an exponentially increasing scale factor on our brane. The proper choice of solution in Case II depends on the initial value of $\dot{b}_o$.

5.3 Case III: $\frac{\kappa^2}{12}\Lambda_{br}^2 - \Lambda_{bulk} \neq 0$ and $\dot{b} \neq 0$

In [2] we also consider the case of additional cosmological constants in the brane and bulk. The results can be found in our longer paper. Not surprisingly we recover the usual cosmological constant driven inflation when we set $\rho = 0$.

6 Sustaining inflation and solving the Horizon Problem

In the previous sections we have found the onset of inflation via superluminal growth of the scale factor. We also know that eventually, once the temperature drops out of the Hagedorn regime, inflation ends. This graceful exit from inflation is an appealing feature of this scenario. However, we cannot calculate in between. Once inflation begins, our calculations break down; our interpretation of small changes in the metric corresponding to small changes in volume no longer holds. At this point we can only speculate what happens in between, and suggest two possibilities.

First, previous authors have argued that strings in a de Sitter background are unstable to fluctuations and that this instability can sustain a period of de Sitter inflation [13, 14, 15]. This phenomenon is well known for strings once they are in the de Sitter-like phase¹. However the missing ingredient that the present study adds is an explanation for how the universe enters a de Sitter like phase in the first place. Our mechanism gets the universe into the locally de Sitter phase, whereupon the mechanism of [13, 14, 15] keeps it there *without* having to assume adiabaticity.

¹We qualify ‘de Sitter’ because the exponential solutions we have do not possess the full O(1,4) de Sitter symmetry.

Second, our 3-brane sits in a bath of branes and bulk strings with ongoing dynamics such as branes smashing into each other and possibly annihilating. The dynamics is likely to keep the system hot. Hence high temperatures near the Hagedorn temperature may be sustained for a long period; inflation continues during this period. We note that the high entropy of our brane $S \sim 10^{88}$ may originate from the dynamics of this hot brane/string bath. Eventually the system cools, drops out of the Hagedorn regime, and inflation ends.

There is a slight difference between our study and the results of [14] which we should comment on. In [14], the density never goes *above* a critical density, ρ'_c . In our case, on the other hand, we must have $\rho > \rho_c \sim 1$ for the calculation to be valid (*i.e.* to be in the Hagedorn regime). The difference arises because [14] was concerned with *stretched* strings and introduced a cut-off in the momentum integral. This ‘coarse graining’ put an artificial upper limit on the amount of string that can be packed into the volume and effectively removed the Hagedorn behaviour. Conversely, the present paper begins in the regime $\rho > \rho_c$ where the calculations of [14] end.

7 Conclusion and discussion

We have studied the possible cosmological implications of the Hagedorn regime of open strings on D-branes in the weak coupling limit. Our main result is that a gas of open strings can exhibit negative pressure leading naturally to a period of power law or exponential inflation. Hagedorn inflation also has a natural exit since any significant cooling can cause a change in the thermodynamics if winding modes become quenched or if the density drops below the critical density, $\rho_c \sim 1$, needed for the entropy of the Hagedorn phase to be dominant. To summarize, open strings on branes in the hot early Hagedorn phase of the universe provide a mechanism to drive a period of inflation, even in the absence of any potential.

Acknowledgements

We thank D. Chung, C. Deffayet, E. Dudas, K. Olive, G. Servant and C. Savoy for discussions. S.A.A. and I.I.K. thank J. Barbón and E. Rabinovici for a previous collaboration and discussions concerning this work. S.A.A. thanks the C.E.A. Saclay for support during this work. K.F. acknowledges support from the Department of Energy through a grant to the University of Michigan. K.F. thanks CERN in Geneva, Switzerland and the Max Planck Institut fuer Physik in Munich, Germany for hospitality during her stay. I.I.K. was supported in part by PPARC rolling grant PPA/G/O/1998/00567, the EC TMR grant FMRX-CT-96-0090 and by the INTAS grant RFBR - 950567.

References

- [1] A. Guth, Phys. Rev. D **23**, 347 (1981)
- [2] S.A. Abel, K. Freese, and I.I Kogan, *JHEP* **0101** (2001) 039
- [3] R. Hagedorn, *Suppl. Nuovo Cimento* **3** (1965) 147; S. Frautschi, Phys. Rev. D **3**, 2821 (1971); R.D. Carlitz, Phys. Rev. D **5**, 3231 (1972)
- [4] K. Huang and S. Weinberg, Phys. Rev. Lett. **25**, 895 (1970); E. Alvarez, Phys. Rev. D **31**, 418 (1985); Nucl. Phys. **B269**, 596 (1986); M. Bowick and L.C.R. Wijewardhana, Phys. Rev. Lett. **54**, 2485 (1985); B. Sundborg, Nucl. Phys. **B254**, 883 (1985); S.N. Tye, Phys. Lett. **158B**, 388 (1985); P. Salomonson and B. Skagerstam, Nucl. Phys. **B268**, 349 (1986); *Physica* **A158** (1989) 499; E. Alvarez and M.A.R. Osorio, Phys. Rev. D **36**, 1175 (1987); D. Mitchell and N. Turok, Phys. Rev. Lett. **58**, 1577 (1987); Nucl. Phys. **B294**, 1138 (1987); I. Antoniadis, J. Ellis and D.V. Nanopoulos, Phys. Lett. **199B**, 402 (1987); M. Axenides, S.D. Ellis and C. Kounnas, Phys. Rev. D **37**, 2964; (1988); I.I. Kogan, JETP. Lett. **45** (1987) 709; B.Sathiapalan, Phys. Rev. D **35**, 3277 (1987); J. Attick and E. Witten, Nucl. Phys. **B310**, 291 (1988); A.A. Abrikosov Jr. and Ya. I. Kogan *Int. J. Mod. Phys.* **A6** (1991) 1501 (submitted 1989), *Sov. Phys. JETP* **69** (1989) 235; R. Brandenberger and C. Vafa, Nucl. Phys. **B316**, 391 (1989); M.J. Bowick and S.B. Giddings, Nucl. Phys. **B325**, 631 (1989); S.B. Giddings, Phys. Lett. **226B**, 55. (1989); B.A. Campbell, N. Kaloper, K.A. Olive, Phys. Lett. **277B**, 265 (1992); S.A. Abel, Nucl. Phys. **B372**, 189 (1992); N. Kaloper, R. Madden, K.A. Olive, Nucl. Phys. **B452**, 677, (1995); M.L. Meana, M.A.R. Osorio and J.P. Penalba, Phys. Lett. **400B**, 275, (1997), Phys. Lett. **408B**, 183 (1997).
- [5] N. Deo, S. Jain and C.-I. Tan, Phys. Lett. **220B**, 125 (1989); Phys. Rev. D **40**, 2646 (1989); N. Deo, S. Jain, O. Narayan and C.-I. Tan, Phys. Rev. D **45**, 3641 (1992)
- [6] D.A. Lowe and L. Thorlacius, Phys. Rev. D **51**, 665 (1995), S. Lee and L. Thorlacius, Phys. Lett. **413B**, 303 (1997).
- [7] J.L.F. Barbón, I.I. Kogan and E. Rabinovici, Nucl. Phys. **B544**, 1999 (104), S.A. Abel, J.L.F. Barbón, I.I. Kogan, and E. Rabinovici, *JHEP* **04** (1999) 015
- [8] M.L. Meana and J.P. Penalba, Nucl. Phys. **B560**, 154 (1999); Phys. Lett. **447B**, 59 (1999); M.A. Vazquez-Mozo, Phys. Rev. D **60**, 106010 (1999); B. Sundborg, hep-th/9908001; S.A. Abel, J.L.F. Barbón, I.I. Kogan, and E. Rabinovici, hep-th/9911004

- [9] For reviews see: J. Polchinski, *TASI* lectures on D-branes, **hep-th/9611050**; “String Theory”, vols 1,2 (CUP) 1998.
- [10] P. Kanti, I. I. Kogan, K. A. Olive, M. Pospelov.
- [11] K. Benakli, Int. J. Mod. Phys. **D8** (1999) 153, A. Lukas, B.A. Ovrut and D. Waldram, **hep-th/9902071**; H.A. Chamblin and H.S. Reall, Nucl. Phys. **B562**, 133 (1999); P. Binétruy, C. Deffayet and D. Langlois, Phys.Lett. **B544** (2002) 183;
- [12] D. Chung and K. Freese, Phys. Rev. D **61**, 023511 (2000).
- [13] R. Brout, F. Englert and E. Gunzig, Ann. Phys. **115** (1978) 78; Gen. Rel. Grav. **10** (1979) 1; R. Brout, F. Englert and P. Spindel, Phys. Rev. Lett. **43**, 417 (1979); Y. Aharanov and A. Casher, Phys. Lett. **166B**, 289 (1986); Y. Aharanov, F. Englert and J. Orloff, Phys. Lett. **199B**, 366 (1987)
- [14] N.Turok, Phys. Rev. Lett. **60**, 549 (1988)
- [15] N. Sanchez and G. Veneziano, Nucl. Phys. **B333** (1990) 253; M. Gasperini, N. Sanchez and G. Veneziano, Int. J. Mod. Phys. **A6** (1991) 3853; Nucl. Phys. **B364** (1991) 365;
- [16] A. Riotto, Phys.Rev. **D61** (2000) 123506.